



**EN** INSTRUKCTIONS FOR USE

Medisorb R-AG  
Resorbable dressing

**CHARACTERISTICS**  
Medisorb R-AG is a sterile, absorbent, non-adhesive, non-occlusive dressing with adhesive borders. It is made of a soft, porous, non-woven fabric. The dressing is designed for use on clean, dry, intact skin. It is suitable for use on wounds, burns, lacerations, and other skin injuries. The dressing is designed to be used for up to 14 days.

**INDICATIONS**  
The dressing is intended for use in the treatment of acute, chronic, and traumatic wounds, lacerations, abrasions, and other skin injuries. It is also suitable for use on burns, lacerations, and other skin injuries. The dressing is designed to be used for up to 14 days.

**CONTRAINDICATIONS**  
The dressing should not be used on infected wounds, wounds with necrotic tissue, or wounds with exposed bone or tendon. It should not be used on wounds that are being treated with systemic antibiotics.

**APPLICATION**  
Before applying the dressing, the wound should be cleaned and dried. The dressing should be applied to the wound surface, ensuring that the adhesive borders are firmly attached to the surrounding skin. The dressing should be changed every 7-14 days.

**PRECAUTIONS**  
The dressing is a sterile product and should be handled with care. It should be stored in its original packaging until ready for use. The dressing should be discarded after use.

**STERILIZATION**  
The dressing is sterilized by gamma irradiation. It is suitable for use on wounds that are being treated with systemic antibiotics.

**STORAGE**  
The dressing should be stored in its original packaging until ready for use. The dressing should be stored at room temperature (15°C to 30°C).

**DATE OF VALIDATION**  
03/2021

**DE** GEBRAUCHSANLEITUNG

Medisorb R-AG  
Resorbierendes Wundnetz

**CHARAKTERISTIKA**  
Medisorb R-AG ist ein steriles, absorbierendes, nicht-adhäsives, nicht-occlusives Wundnetz mit klebenden Rändern. Es besteht aus einem weichen, porösen, nichtgewebten Material. Das Wundnetz ist für den Einsatz auf saubere, trockene, intakte Haut vorgesehen. Es eignet sich für die Behandlung von Wunden, Schürfwunden, Schnittwunden und anderen Hautverletzungen. Das Wundnetz ist für einen Einsatz von bis zu 14 Tagen vorgesehen.

**ANWENDUNG**  
Vor dem Aufbringen des Wundnetzes sollte die Wunde gereinigt und getrocknet werden. Das Wundnetz sollte auf die Wundfläche aufgebracht werden, wobei die klebenden Ränder fest an der umliegenden Haut befestigt werden. Das Wundnetz sollte alle 7 bis 14 Tage gewechselt werden.

**VORSICHTSMAßNAHMEN**  
Das Wundnetz ist ein steriles Produkt und sollte sorgfältig gehandhabt werden. Es sollte in der originalen Verpackung bis zur Verwendung aufbewahrt werden. Das Wundnetz sollte nach dem Gebrauch entsorgt werden.

**STERILISATION**  
Das Wundnetz wird durch Gammastrahlung sterilisiert. Es ist für die Behandlung von Wunden geeignet, die mit systemischen Antibiotika behandelt werden.

**LAGERUNG**  
Das Wundnetz sollte in der originalen Verpackung bis zur Verwendung aufbewahrt werden. Das Wundnetz sollte bei Raumtemperatur (15°C bis 30°C) gelagert werden.

**VALIDATIONSDATUM**  
03/2021

**FR** MODE D'EMPLOI

Medisorb R-AG  
Pansement résorbable

**CHARACTÉRISTIQUES**  
Le pansement est stérile, absorbant, non adhésif, non occlusif. Il est composé d'un tissu non tissé doux et poreux. Le pansement est conçu pour être utilisé sur une peau propre, sèche et intacte. Il est adapté pour traiter les plaies, les brûlures, les lésions et d'autres blessures cutanées. Le pansement est conçu pour être utilisé pendant jusqu'à 14 jours.

**INDICATIONS**  
Le pansement est destiné à être utilisé pour le traitement des plaies aiguës, chroniques et traumatiques, des brûlures, des lésions et d'autres blessures cutanées. Il est également adapté pour être utilisé sur des brûlures, des lésions et d'autres blessures cutanées.

**CONTRA-INDICATIONS**  
Le pansement ne doit pas être utilisé sur des plaies infectées, des plaies à tissu nécrotique ou des plaies avec des os ou des tendons exposés. Il ne doit pas être utilisé sur des plaies traitées avec des antibiotiques systémiques.

**MODE D'EMPLOI**  
Avant d'appliquer le pansement, nettoyez et séchez soigneusement la plaie. Appliquez le pansement sur la surface de la plaie, en veillant à ce que les bords adhésifs soient bien collés à la peau saine environnante. Changez le pansement tous les 7 à 14 jours.

**PRÉCAUTIONS**  
Le pansement est un produit stérile et doit être manipulé avec soin. Il doit être conservé dans son emballage d'origine jusqu'à son utilisation. Le pansement doit être jeté après utilisation.

**STÉRILISATION**  
Le pansement est stérilisé par irradiation gamma. Il est adapté pour être utilisé sur des plaies traitées avec des antibiotiques systémiques.

**STOCKAGE**  
Le pansement doit être conservé dans son emballage d'origine jusqu'à son utilisation. Le pansement doit être conservé à température ambiante (15°C à 30°C).

**DATE DE VALIDATION**  
03/2021

**NL** INSTRUCTIES VOOR GEBRUIK

Medisorb R-AG  
Resorbabel dressing

**CHARACTERISTIEKEN**  
Medisorb R-AG is een steriel, absorberend, niet-adhese, niet-occlusief wondverband met kleefranden. Het is vervaardigd uit een zacht, poreus, niet-geweven materiaal. Het wondverband is bedoeld voor gebruik op schone, droge, intacte huid. Het is geschikt voor het behandelen van wonden, brandwonden, snijwonden en andere huidwonden. Het wondverband is bedoeld voor gebruik tot maximaal 14 dagen.

**INDICATIES**  
Het wondverband is bedoeld voor de behandeling van acuute, chronische, traumatische wonden, brandwonden, snijwonden en andere huidwonden. Het is ook geschikt voor gebruik op brandwonden, snijwonden en andere huidwonden.

**CONTRA-INDICATIES**  
Het wondverband mag niet worden gebruikt op geïnfecteerde wonden, wonden met necrotisch weefsel of wonden met blootliggende botten of tendons. Het mag niet worden gebruikt op wonden die worden behandeld met systemische antibiotica.

**AFGEVOELD**  
Voordat u het wondverband aanbreft, moet u de wond grondig reinigen en drogen. Het wondverband moet op de wondoppervlakte worden aangebracht, waarbij de kleefranden stevig aan de omliggende huid worden vastgezet. Het wondverband moet elke 7 tot 14 dagen worden vervangen.

**OPMERKINGEN**  
Het wondverband is een steriel product en moet zorgvuldig worden behandeld. Het moet in de originele verpakking tot gebruik worden bewaard. Het wondverband moet na gebruik worden weggegooid.

**STERILISATIE**  
Het wondverband wordt gesteriliseerd door gammastraling. Het is geschikt voor gebruik op wonden die worden behandeld met systemische antibiotica.

**OPBERGING**  
Het wondverband moet in de originele verpakking tot gebruik worden bewaard. Het wondverband moet bij kamertemperatuur (15°C tot 30°C) worden opgeslagen.

**VALIDATIEDATUM**  
03/2021

**PT** INSTRUKÇÕES FOR USE

Medisorb R-AG  
Pano resorçavel

**CARACTERÍSTICAS**  
O Medisorb R-AG é um produto estéril, absorvente, não adesivo, não oclusivo, com bordas adesivas. É feito de um tecido não tecido macio e poroso. O produto é destinado para uso em pele limpa, seca e íntegra. É adequado para tratar feridas, queimaduras, cortes e outras lesões cutâneas. O produto é destinado para uso por até 14 dias.

**INDICAÇÕES**  
O produto deve ser usado no tratamento de feridas agudas, crônicas e traumáticas, queimaduras, cortes e outras lesões cutâneas. Também é adequado para uso em queimaduras, cortes e outras lesões cutâneas.

**CONTRA-INDICAÇÕES**  
O produto não deve ser usado em feridas infectadas, feridas com tecido necrótico ou feridas com osso ou tendão expostos. Não deve ser usado em feridas tratadas com antibióticos sistêmicos.

**APLICAÇÃO**  
Antes de aplicar o produto, a ferida deve ser limpa e seca. O produto deve ser aplicado sobre a superfície da ferida, garantindo que as bordas adesivas estejam firmemente coladas à pele saudável adjacente. O produto deve ser trocado a cada 7 a 14 dias.

**PRECAUÇÕES**  
O produto é um produto estéril e deve ser manuseado com cuidado. Deve ser armazenado em sua embalagem original até o uso. O produto deve ser descartado após o uso.

**ESTERILIZAÇÃO**  
O produto é esterilizado por irradiação gama. É adequado para uso em feridas tratadas com antibióticos sistêmicos.

**ARMAZENAMENTO**  
O produto deve ser armazenado em sua embalagem original até o uso. O produto deve ser armazenado em temperatura ambiente (15°C a 30°C).

**DATA DE VALIDAÇÃO**  
03/2021

**HU** HASZNÁLTÁJUTÓMATÓ

Medisorb R-AG  
Reszoráló kötség

**JELLEMZŐK**  
A Medisorb R-AG egy steril, felszívó, nem tapadós, nem oltató kötszék ragasztó széllel. A kötszék szövet puha és póros. A kötszék a tisztán, száraz, érintetlen bőrre való használatra készült. Alkalmazható a sebek, égések, vágások, és egyéb bőrelváltozások kezelésére. A kötszék használható legfeljebb 14 napig.

**ALKALMAZÁS**  
A kötszék alkalmazása előtt a sebet alaposan tisztítani kell. A kötszék a sebre kell lenyújtani, ügyelve arra, hogy a ragasztó szél szorosan tapadjon a környező bőrre. A kötszék minden 7-14 naponta cserélendő.

**FIGYELMEZTETÉSEK**  
A kötszék nem alkalmazható fertőzött sebekre, nem nekrotikus szövetekkel borított sebekre, valamint olyan sebekre, amelyeknél kiemelkednek az erek vagy a idegek. Ne használja a kötszék a szisztémás antibiotikumokkal történő kezeléssel együtt.

**FIGYELMEZTETÉSEK**  
A kötszék egy steril termék, ezért gondosan kell kezelni. A kötszék eredeti csomagolásában kell tartani, amíg nem használja. A kötszék használatát követően meg kell semmisíteni.

**ESTERILIZÁCIÓ**  
A kötszék gamma sugárzással sterilizálva van. Alkalmazható olyan sebek kezelésére, amelyeknél szisztémás antibiotikumokat alkalmaznak.

**TÁROLÁS**  
A kötszék eredeti csomagolásában kell tartani, amíg nem használja. A kötszék tárolási hőmérséklete szobahőmérsékleten (15°C-30°C) lehet.

**ÉRTÉKELÉS DÁTUMA**  
03/2021

**LOT**

Medisorb R-AG

**CHARACTERISTICS**  
Medisorb R-AG is a sterile, absorbent, non-adhesive, non-occlusive dressing with adhesive borders. It is made of a soft, porous, non-woven fabric. The dressing is designed for use on clean, dry, intact skin. It is suitable for use on wounds, burns, lacerations, and other skin injuries. The dressing is designed to be used for up to 14 days.

**INDICATIONS**  
The dressing is intended for use in the treatment of acute, chronic, and traumatic wounds, lacerations, abrasions, and other skin injuries. It is also suitable for use on burns, lacerations, and other skin injuries. The dressing is designed to be used for up to 14 days.

**CONTRAINDICATIONS**  
The dressing should not be used on infected wounds, wounds with necrotic tissue, or wounds with exposed bone or tendon. It should not be used on wounds that are being treated with systemic antibiotics.

**APPLICATION**  
Before applying the dressing, the wound should be cleaned and dried. The dressing should be applied to the wound surface, ensuring that the adhesive borders are firmly attached to the surrounding skin. The dressing should be changed every 7-14 days.

**PRECAUTIONS**  
The dressing is a sterile product and should be handled with care. It should be stored in its original packaging until ready for use. The dressing should be discarded after use.

**STERILIZATION**  
The dressing is sterilized by gamma irradiation. It is suitable for use on wounds that are being treated with systemic antibiotics.

**STORAGE**  
The dressing should be stored in its original packaging until ready for use. The dressing should be stored at room temperature (15°C to 30°C).

**DATE OF VALIDATION**  
03/2021

**LOT**

Medisorb R-AG

**CHARACTERISTICS**  
Medisorb R-AG is a sterile, absorbent, non-adhesive, non-occlusive dressing with adhesive borders. It is made of a soft, porous, non-woven fabric. The dressing is designed for use on clean, dry, intact skin. It is suitable for use on wounds, burns, lacerations, and other skin injuries. The dressing is designed to be used for up to 14 days.

**INDICATIONS**  
The dressing is intended for use in the treatment of acute, chronic, and traumatic wounds, lacerations, abrasions, and other skin injuries. It is also suitable for use on burns, lacerations, and other skin injuries. The dressing is designed to be used for up to 14 days.

**CONTRAINDICATIONS**  
The dressing should not be used on infected wounds, wounds with necrotic tissue, or wounds with exposed bone or tendon. It should not be used on wounds that are being treated with systemic antibiotics.

**APPLICATION**  
Before applying the dressing, the wound should be cleaned and dried. The dressing should be applied to the wound surface, ensuring that the adhesive borders are firmly attached to the surrounding skin. The dressing should be changed every 7-14 days.

**PRECAUTIONS**  
The dressing is a sterile product and should be handled with care. It should be stored in its original packaging until ready for use. The dressing should be discarded after use.

**STERILIZATION**  
The dressing is sterilized by gamma irradiation. It is suitable for use on wounds that are being treated with systemic antibiotics.

**STORAGE**  
The dressing should be stored in its original packaging until ready for use. The dressing should be stored at room temperature (15°C to 30°C).

**DATE OF VALIDATION**  
03/2021

**LOT**

Medisorb R-AG

**CHARACTERISTICS**  
Medisorb R-AG is a sterile, absorbent, non-adhesive, non-occlusive dressing with adhesive borders. It is made of a soft, porous, non-woven fabric. The dressing is designed for use on clean, dry, intact skin. It is suitable for use on wounds, burns, lacerations, and other skin injuries. The dressing is designed to be used for up to 14 days.

**INDICATIONS**  
The dressing is intended for use in the treatment of acute, chronic, and traumatic wounds, lacerations, abrasions, and other skin injuries. It is also suitable for use on burns, lacerations, and other skin injuries. The dressing is designed to be used for up to 14 days.

**CONTRAINDICATIONS**  
The dressing should not be used on infected wounds, wounds with necrotic tissue, or wounds with exposed bone or tendon. It should not be used on wounds that are being treated with systemic antibiotics.

**APPLICATION**  
Before applying the dressing, the wound should be cleaned and dried. The dressing should be applied to the wound surface, ensuring that the adhesive borders are firmly attached to the surrounding skin. The dressing should be changed every 7-14 days.

**PRECAUTIONS**  
The dressing is a sterile product and should be handled with care. It should be stored in its original packaging until ready for use. The dressing should be discarded after use.

**STERILIZATION**  
The dressing is sterilized by gamma irradiation. It is suitable for use on wounds that are being treated with systemic antibiotics.

**STORAGE**  
The dressing should be stored in its original packaging until ready for use. The dressing should be stored at room temperature (15°C to 30°C).

**DATE OF VALIDATION**  
03/2021

**LOT**

Medisorb R-AG

**CHARACTERISTICS**  
Medisorb R-AG is a sterile, absorbent, non-adhesive, non-occlusive dressing with adhesive borders. It is made of a soft, porous, non-woven fabric. The dressing is designed for use on clean, dry, intact skin. It is suitable for use on wounds, burns, lacerations, and other skin injuries. The dressing is designed to be used for up to 14 days.

**INDICATIONS**  
The dressing is intended for use in the treatment of acute, chronic, and traumatic wounds, lacerations, abrasions, and other skin injuries. It is also suitable for use on burns, lacerations, and other skin injuries. The dressing is designed to be used for up to 14 days.

**CONTRAINDICATIONS**  
The dressing should not be used on infected wounds, wounds with necrotic tissue, or wounds with exposed bone or tendon. It should not be used on wounds that are being treated with systemic antibiotics.

**APPLICATION**  
Before applying the dressing, the wound should be cleaned and dried. The dressing should be applied to the wound surface, ensuring that the adhesive borders are firmly attached to the surrounding skin. The dressing should be changed every 7-14 days.

**PRECAUTIONS**  
The dressing is a sterile product and should be handled with care. It should be stored in its original packaging until ready for use. The dressing should be discarded after use.

**STERILIZATION**  
The dressing is sterilized by gamma irradiation. It is suitable for use on wounds that are being treated with systemic antibiotics.

**STORAGE**  
The dressing should be stored in its original packaging until ready for use. The dressing should be stored at room temperature (15°C to 30°C).

**DATE OF VALIDATION**  
03/2021

**LOT**

Medisorb R-AG

**CHARACTERISTICS**  
Medisorb R-AG is a sterile, absorbent, non-adhesive, non-occlusive dressing with adhesive borders. It is made of a soft, porous, non-woven fabric. The dressing is designed for use on clean, dry, intact skin. It is suitable for use on wounds, burns, lacerations, and other skin injuries. The dressing is designed to be used for up to 14 days.

**INDICATIONS**  
The dressing is intended for use in the treatment of acute, chronic, and traumatic wounds, lacerations, abrasions, and other skin injuries. It is also suitable for use on burns, lacerations, and other skin injuries. The dressing is designed to be used for up to 14 days.

**CONTRAINDICATIONS**  
The dressing should not be used on infected wounds, wounds with necrotic tissue, or wounds with exposed bone or tendon. It should not be used on wounds that are being treated with systemic antibiotics.

**APPLICATION**  
Before applying the dressing, the wound should be cleaned and dried. The dressing should be applied to the wound surface, ensuring that the adhesive borders are firmly attached to the surrounding skin. The dressing should be changed every 7-14 days.

**PRECAUTIONS**  
The dressing is a sterile product and should be handled with care. It should be stored in its original packaging until ready for use. The dressing should be discarded after use.

**STERILIZATION**  
The dressing is sterilized by gamma irradiation. It is suitable for use on wounds that are being treated with systemic antibiotics.

**STORAGE**  
The dressing should be stored in its original packaging until ready for use. The dressing should be stored at room temperature (15°C to 30°C).

**DATE OF VALIDATION**  
03/2021

**LOT**

Medisorb R-AG

**CHARACTERISTICS**  
Medisorb R-AG is a sterile, absorbent, non-adhesive, non-occlusive dressing with adhesive borders. It is made of a soft, porous, non-woven fabric. The dressing is designed for use on clean, dry, intact skin. It is suitable for use on wounds, burns, lacerations, and other skin injuries. The dressing is designed to be used for up to 14 days.

**INDICATIONS**  
The dressing is intended for use in the treatment of acute, chronic, and traumatic wounds, lacerations, abrasions, and other skin injuries. It is also suitable for use on burns, lacerations, and other skin injuries. The dressing is designed to be used for up to 14 days.

**CONTRAINDICATIONS**  
The dressing should not be used on infected wounds, wounds with necrotic tissue, or wounds with exposed bone or tendon. It should not be used on wounds that are being treated with systemic antibiotics.

**APPLICATION**  
Before applying the dressing, the wound should be cleaned and dried. The dressing should be applied to the wound surface, ensuring that the adhesive borders are firmly attached to the surrounding skin. The dressing should be changed every 7-14 days.

**PRECAUTIONS**  
The dressing is a sterile product and should be handled with care. It should be stored in its original packaging until ready for use. The dressing should be discarded after use.

**STERILIZATION**  
The dressing is sterilized by gamma irradiation. It is suitable for use on wounds that are being treated with systemic antibiotics.

**STORAGE**  
The dressing should be stored in its original packaging until ready for use. The dressing should be stored at room temperature (15°C to 30°C).

**DATE OF VALIDATION**  
03/2021

**LOT**

Medisorb R-AG

**CHARACTERISTICS**  
Medisorb R-AG is a sterile, absorbent, non-adhesive, non-occlusive dressing with adhesive borders. It is made of a soft, porous, non-woven fabric. The dressing is designed for use on clean, dry, intact skin. It is suitable for use on wounds, burns, lacerations, and other skin injuries. The dressing is designed to be used for up to 14 days.

**INDICATIONS**  
The dressing is intended for use in the treatment of acute, chronic, and traumatic wounds, lacerations, abrasions, and other skin injuries. It is also suitable for use on burns, lacerations, and other skin injuries. The dressing is designed to be used for up to 14 days.

**CONTRAINDICATIONS**  
The dressing should not be used on infected wounds, wounds with necrotic tissue, or wounds with exposed bone or tendon. It should not be used on wounds that are being treated with systemic antibiotics.

**APPLICATION**  
Before applying the dressing, the wound should be cleaned and dried. The dressing should be applied to the wound surface, ensuring that the adhesive borders are firmly attached to the surrounding skin. The dressing should be changed every 7-14 days.

**PRECAUTIONS**  
The dressing is a sterile product and should be handled with care. It should be stored in its original packaging until ready for use. The dressing should be discarded after use.

**STERILIZATION**  
The dressing is sterilized by gamma irradiation. It is suitable for use on wounds that are being treated with systemic antibiotics.

**STORAGE**  
The dressing should be stored in its original packaging until ready for use. The dressing should be stored at room temperature (15°C to 30°C).

**DATE OF VALIDATION**  
03/2021

**LOT**

Medisorb R-AG

**CHARACTERISTICS**  
Medisorb R-AG is a sterile, absorbent, non-adhesive, non-occlusive dressing with adhesive borders. It is made of a soft, porous, non-woven fabric. The dressing is designed for use on clean, dry, intact skin. It is suitable for use on wounds, burns, lacerations, and other skin injuries. The dressing is designed to be used for up to 14 days.

**INDICATIONS**  
The dressing is intended for use in the treatment of acute, chronic, and traumatic wounds, lacerations, abrasions, and other skin injuries. It is also suitable for use on burns, lacerations, and other skin injuries. The dressing is designed to be used for up to 14 days.

**CONTRAINDICATIONS**  
The dressing should not be used on infected wounds, wounds with necrotic tissue, or wounds with exposed bone or tendon. It should not be used on wounds that are being treated with systemic antibiotics.

**APPLICATION**  
Before applying the dressing, the wound should be cleaned and dried. The dressing should be applied to the wound surface, ensuring that the adhesive borders are firmly attached to the surrounding skin. The dressing should be changed every 7-14 days.

**PRECAUTIONS**  
The dressing is a sterile product and should be handled with care. It should be stored in its original packaging until ready for use. The dressing should be discarded after use.

**STERILIZATION**  
The dressing is sterilized by gamma irradiation. It is suitable for use on wounds that are being treated with systemic antibiotics.

**STORAGE**  
The dressing should be stored in its original packaging until ready for use. The dressing should be stored at room temperature (15°C to 30°C).

**DATE OF VALIDATION**  
03/2021

**LOT**

Medisorb R-AG

**CHARACTERISTICS**  
Medisorb R-AG is a sterile, absorbent, non-adhesive, non-occlusive dressing with adhesive borders. It is made of a soft, porous, non-woven fabric. The dressing is designed for use on clean, dry, intact skin. It is suitable for use on wounds, burns, lacerations, and other skin injuries. The dressing is designed to be used for up to 14 days.

**INDICATIONS**  
The dressing is intended for use in the treatment of acute, chronic, and traumatic wounds, lacerations, abrasions, and other skin injuries. It is also suitable for use on burns, lacerations, and other skin injuries. The dressing is designed to be used for up to 14 days.

**CONTRAINDICATIONS**  
The dressing should not be used on infected wounds, wounds with necrotic tissue, or wounds with exposed bone or tendon. It should not be used on wounds that are being treated with systemic antibiotics.

**APPLICATION**  
Before applying the dressing, the wound should be cleaned and dried. The dressing should be applied to the wound surface, ensuring that the adhesive borders are firmly attached to the surrounding skin. The dressing should be changed every 7-14 days.

**PRECAUTIONS**  
The dressing is a sterile product and should be handled with care. It should be stored in its original packaging until ready for use. The dressing should be discarded after use.

**STERILIZATION**  
The dressing is sterilized by gamma irradiation. It is suitable for use on wounds that are being treated with systemic antibiotics.

**STORAGE**  
The dressing should be stored in its original packaging until ready for use. The dressing should be stored at room temperature (15°C to 30°C).

**DATE OF VALIDATION**  
03/2021

**LOT**

Medisorb R-AG

**CHARACTERISTICS**  
Medisorb R-AG is a sterile, absorbent, non-adhesive, non-occlusive dressing with adhesive borders. It is made of a soft, porous, non-woven fabric. The dressing is designed for use on clean, dry, intact skin. It is suitable for use on wounds, burns, lacerations, and other skin injuries. The dressing is designed to be used for up to 14 days.

**INDICATIONS**  
The dressing is intended for use in the treatment of acute, chronic, and traumatic wounds, lacerations, abrasions, and other skin injuries. It is also suitable for use on burns, lacerations, and other skin injuries. The dressing is designed to be used for up to 14 days.

**CONTRAINDICATIONS**  
The dressing should not be used on infected wounds, wounds with necrotic tissue, or wounds with exposed bone or tendon. It should not be used on wounds that are being treated with systemic antibiotics.

**APPLICATION**  
Before applying the dressing, the wound should be cleaned and dried. The dressing should be applied to the wound surface, ensuring that the adhesive borders are firmly attached to the surrounding skin. The dressing should be changed every 7-14 days.

**PRECAUTIONS**  
The dressing is a sterile product and should be handled with care. It should be stored in its original packaging until ready for use. The dressing should be discarded after use.

**STERILIZATION**  
The dressing is sterilized by gamma irradiation. It is suitable for use on wounds that are being treated with systemic antibiotics.

**STORAGE**  
The dressing should be stored in its original packaging until ready for use. The dressing should be stored at room temperature (